



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## NOTES ON STAR DIAMETERS:

I. POSSIBLE VARIATIONS IN THE DIAMETER OF  $\alpha$  ORIONIS

Further measures have been made on  $\alpha$  *Orionis* with the 20-foot interferometer attached to the 100-inch reflector, with the beam extending east and west. On the nights of October 14 and 15, 1922, the visibility curve was found to cross the axis at 14 feet, corresponding to an angular diameter of  $0''.034$ . On the first night the seeing was excellent; on the second night it



was poor, but there were enough moments of good seeing to obtain an excellent curve. The readings for minimum diameter last year were  $8\frac{1}{2}$  feet, equivalent to a diameter of  $0''.054$ . These measures were made in September and November, 1921, and the seeing was the same as on October 15 of this year. For the year 1920, we have the measures made in December with good seeing, giving a disappearance of the fringes at ten feet,

equivalent to an angular diameter of  $0''.047$ . It is believed these measures are not more than ten per cent in error.

Two suggestions are offered to explain these different results: First, the seeing "background" may be better this year than in the two previous years. This does not seem to be upheld by readings made on the zero images and on other stars. The ordinary seeing was, in general, poor throughout the year, but there were times of good interferometer seeing which were not far different from those of this year. Second, there may be some variation in the effective diameter of the star itself. It is well known that  $\alpha$  *Orionis* varies in brightness, and on the accompanying chart I have shown a rough curve of Osthoff's measures<sup>1</sup> of its visual brightness compared with  $\beta$  *Orionis* and  $\alpha$  *Tauri*. Professor Barnard has generously placed at my disposal his unpublished naked-eye comparisons of  $\alpha$  *Orionis* with  $\alpha$  *Tauri*, and upon plotting these we find the curve not to differ greatly from Osthoff's. I have also inserted Lunt's plotting of Bottlinger's radial velocity curve,<sup>2</sup> extending it for a period of 12 years. Plotting the dates of interferometer observations of minimum diameter upon the radial velocity curve, we have found that the ordinates of the curve are roughly proportional to the mirror separations. Lunt believes that Bottlinger's curve needs to be corrected and that the maximum and minimum occur earlier than shown. From the present data it appears that some relationship may exist between radial velocity, visual brightness and angular diameter, the phase relation as yet being uncertain. Additional interferometer measurements are necessary to determine whether this connection is real or accidental. The diameters given are calculated from the formula for a uniform disk,  $a = 1.22\lambda/d$ . It is not at all certain that the factor 1.22 should be used throughout. It would aid greatly in solving the problem to have further precise observations of the visual magnitude, bolometric intensity, and radial velocity, and also to note particularly whether there may not be extremely small changes in type of spectrum.

FRANCIS G. PEASE

Mount Wilson Observatory, November 6, 1922.

<sup>1</sup>*Die Veränderlichkeit von  $\alpha$  Orionis*, Von H. Osthoff. *A. N.* 5172, 187.

<sup>2</sup>*On the Orbits of the Spectroscopic Binaries  $\alpha$  Orionis and  $\alpha$  Scorpii*. Joseph